

Nonstationary Transport by Internal White Noise and "Localization" in Ratchets

Nikolai Brilliantov^{1,2} and Peter Strizhak^{1,3}

¹ *Department of Chemistry, University of Toronto, Toronto, Canada M5S 3H6*

² *Physics Department, Moscow State University, Moscow 119899, Russia*

³ *Institute of Physical Chemistry, National Academy of Sciences of Ukraine, pr.Nauki, 31, Kiev 252038, Ukraine*

We revealed a right-left asymmetry of the inter-well mean first passage times for the Brownian particles in a ratchet potential under internal white noise. We showed analytically and numerically that this asymmetry gives rise to the following phenomena: (i) nonstationary transport; (ii) localization or ultraslow (Sinai) diffusion for ratchets with a disorder in the asymmetry parameter of the potential; (iii) equilibrium "drift of labels" in circular ratchets.

PACS numbers: 82.20.Mj, 05.40.+j

It was demonstrated recently, that nonequilibrium fluctuations may support a stationary flux of the Brownian particles in a periodic potential with broken symmetry (ratchet). This fluctuation-induced transport has been of growing interest after Magnasco suggestion to apply a toy "ratchet and pawl" Feynman's model to riddle a "mystery" of the directed transport in cells [1]. In subsequent studies [2–7], some related models were proposed to explain the nature of molecular motors [2,4], to develop a novel mass separation technique [7], to design the Brownian engine with a negative resistance [8], or to speculate how the ion channel junctions operate [6]. According to the second law, no stationary transport occurs under equilibrium fluctuations. To obtain a stationary motion, one has to break somehow the detailed balance. A possible way is to impose an additional colored noise [1–3,9–11]. One can also couple the system with nonequilibrium bath [5], with external (e.g. chemical) degrees of freedom [2,4,8], or with a deterministic chaotic process [12].

In the present study we consider the motion of the Brownian particles in a ratchet potential (Fig.1a) under an equilibrium white noise. We found a right-left asymmetry for the mean first passage times (MFPT) of the interwell transitions. This property of the MFPT entails the following phenomena: (i) nonstationary transport of the Brownian particles; (ii) ultraslow (Sinai) diffusion or "localization" in ratchets with a disorder (Fig.1b.); (iii) equilibrium "drift of labels" in circular ratchets. The directed nonstationary transport of the ensemble of the Brownian particles occurs as its peculiar route towards equilibrium [13]. If the process is interrupted at some stage (e.g. removing a particle) a net transport is obtained. This scenario of the nonstationary transport seems to be realistic for biological systems.

Nothing is perfectly periodic in nature, and any periodic structure has imperfections and distortions. Sometimes even weak disorder drastically changes a kinetic behavior of the system [14]. If the ratchet potential is symmetric in average, but the asymmetry parameter, $\alpha = L_-/L_0$ is stochastically distributed around $\alpha = 1/2$ (Fig.1), the mean displacement of the particles is zero.

Surprisingly, the mean square displacement behaves as $\sim [\log(t)]^4$ at large times t , contrary to $\sim t$ dependence for the usual diffusion. At $t \gg 1$ $[\log(t)]^4/t \rightarrow 0$, so the particle is, in fact, localized on the length scale of the ordinary diffusion. The "localization" may be effectively used in biological systems, or in technology, instead of trapping molecules in a deep potential well by forming a chemical bond. It costs much less energy to release the particle localized by the disordered ratchets, rather than waste energy, breaking the chemical bond.

The directed nonstationary transport and ultraslow diffusion concern the Brownian motion on the infinite line. The other interesting phenomenon occurs in circular ratchets. From one hand, the equilibrium flux in a ring is zero, from the other, the interwell MFPT for transitions to the right is not equal to the MFPT of the left transitions. As a result a "drift of labels" may be detected, provided that the particles may be labeled and their paths may be traced. The "drift of labels" is not a thermodynamical flux and the detailed balance at equilibrium is preserved.

To address these problems we consider the Brownian particle in a periodic ratchet potential (Fig.1a), subjected to the white noise, $\xi(t)$, $\langle \xi(t)\xi(t') \rangle = 2D\delta(t-t')$. Here $D = kT/\gamma$ is the diffusion, and γ is the friction coefficient. The Langevin equation for the overdamped case reads:

$$\dot{x} = f(x) + \xi(t) \quad (1)$$

with $f(x) = -\gamma^{-1}\partial U/\partial x$, and $U(x)$ being the ratchet potential. The corresponding Fokker-Plank equation is

$$\partial_t P = -\partial_x (fP) + D\partial_{xx}P \quad (2)$$

The Brownian motion in the piecewise linear potential may be effectively studied in terms of the MFPT [15]. The escape time from a starting point, x_0 , in the well is given as a solution of the equation [16]:

$$D \frac{d^2 T(x_0)}{dx_0^2} - \gamma^{-1} \frac{dU}{dx_0} \frac{dT(x_0)}{dx_0} + 1 = 0 \quad (3)$$

The final point is either L_+ , if the particle escapes to the right, or $-L_-$, if it overcomes the barrier in the opposite direction. The averaging of the escape time over x_0 gives T_+ and T_- , the MFPT for the right and for the left transitions respectively. To find T_+ , we use the continuity of $T'(x_0)$, and boundary conditions $T(L_+) = 0$, $T'(-L_-) = 0$, which are self-consistent at $\tau_{\pm}/T_{\pm} \ll 1$. Here $\tau_{\pm} = \gamma L_{\pm}/kTa_{\pm}$ are the *intrawell* relaxation times, and $a_{\pm} = U_0/kTL_{\pm}$ characterize the force on the slopes. The boundary conditions are reversed to obtain T_- . We consider the case when $\exp(U_0/kT) \gg 1$ and $\tau_{\pm}/T_{\pm} \ll 1$, i.e., the particle has enough time to thermalize inside the well before it jumps over the barrier. Straightforward analysis shows that the solution of Eq.(3) is a sum of few terms [17]. The leading term is independent on x_0 , and is of the order of $\exp(U_0/kT) \gg 1$. The other terms [17] may be neglected everywhere except the regions in the very vicinity of the tops. If the distribution inside the well is sharply peaked around the bottom, these regions do not noticeably contribute to the MFPT. Particularly, this is true for the equilibrium distribution if $U_0/kT \geq 0.1$. Finally, we obtain

$$T_{\pm} = a_{\pm}^{-1} e^{U_0/kT} / Da_0 \quad (4)$$

where $a_0^{-1} = a_+^{-1} + a_-^{-1}$. This equation shows that $T_+ \neq T_-$ for the potential with asymmetry. It is worth noting that the left-right asymmetry of the MFPT for the ratchet under the white noise occurs as the preexponential factor. For the colored noise this factor is overshadowed by the asymmetry that appears in the exponent [10]. The left-right asymmetry of the MFPT described by Eq.(4) was confirmed in our numerical studies performed for the equilibrium intrawell distribution. To get the required white noise process (which is effectively colored due to the finite time step) we decreased the time step until the results became insensitive to it further decrease in a range of two decades (the time step varied from $2 \cdot 10^{-4}$ to 10^{-6} , and was roughly two orders of magnitude smaller, then that, used in [18]). We found that the simplest Euler method (see e.g. [18]) is quite effective for the piecewise linear potential.

As a consequence of the left-right asymmetry in the MFPT, the directed motion appears before the system reaches the stationary probability distribution, $P_{eq}(x) \propto \exp(-U(x)/kT)$, which is a current free, if no density gradients present in the system. Thus, any initially localized distribution, say $P(x, t; x_0, t_0) = \delta(x - x_0)(t - t_0)$, evolves to the $P_{eq}(x)$, defined on the ratchet, with left and right boundary $-L_{\infty}$, $L_{\infty} \gg L_0$. In the sequence of the time scales, that characterize the Brownian motion in the overdamped case, the shortest time scale corresponds to the intrawell relaxation time $\tau_{\pm}$. The next time scale is related to the interwells transition times, $T_{\pm}$. Finally, $T_{\infty} \sim (L_{\infty}/L_0)T_{\pm}$ defines the global relaxation time. Under condition $\exp(U_0/kT) \gg 1$, the following time-scales hierarchy holds, $\tau_{\pm} \ll T_{\pm} \ll T_{\infty}$, and the nonstationary transport takes place for $T_{\pm} \ll t \ll T_{\infty}$.

To describe the nonstationary transport we introduce the population of the k -th well as $n_k(t) = \int_{L_0 k}^{L_0(k+1)} P(x, t) dx$, and write the kinetic equation:

$$\dot{n}_k = W_- n_{k+1} + W_+ n_{k-1} - (W_- + W_+) n_k \quad (5)$$

where $W_{\pm}$ give transition rates, which may be estimated as $W_{\pm} \simeq T_{\pm}^{-1}$ [19]. Performing transformation from n_k in the discrete space to $n(x)$ in continuous space coordinate, we obtain the coarse-grained equation:

$$\partial_t n(x, t) \simeq -V \partial_x n(x, t) + \tilde{D} \partial_{xx} n(x, t) \quad (6)$$

with $V = L_0(T_+^{-1} - T_-^{-1})$ and $\tilde{D} = \frac{1}{2} L_0^2(T_+^{-1} + T_-^{-1})$. More rigorous derivation of the coarse-grained equation, based on the separation of different time-scales [20] yields the same result for $t \gg T_{\pm}$.

Eq.(6) shows that the ensemble of Brownian particles moves with a constant velocity V , which increases with enhancing asymmetry of the potential as $V = (U_0/kT)^2 e^{-(U_0/kT)} (D/L_0) (\frac{1}{1-\alpha} - \frac{1}{\alpha})$ (Fig.2). The increase of the distribution width with time is described by the renormalized diffusion coefficient $\tilde{D}$.

We apply the coarse-grained description for the case of the external field $U_{ex}(x) = -Ex$. Calculations of the MFPT for the small fields, $EL_0/U_0 \ll 1$, yield Eq.(4), with $a_{\pm}$ substituted by $a_{\pm} \pm E/kT$, and U_0/kT substituted by $(U_0 \pm EL_{\pm})/kT$. In the linear approximation for E , the average velocity reads:

$$V_E = V + (\tilde{D}/kT)E \quad (7)$$

where V and $\tilde{D}$ are the field-free values. The top insert in Fig.2 depicts dependence of the average velocity upon the external force. The particles move uphill, if the field does not exceed the critical value, $E_c = (2kT/L_0)(1 - 2\alpha)$.

The kinetic behavior of the system in the vicinity of the critical field, i.e. near $V_E = 0$ drastically changes under (even weak) disorder. The disorder may be either in the values of E , or in the values of $T_{\pm}$, i.e. in the asymmetry parameter α . For such a "quenched disorder" [14], one has stochastically distributed transition probability rates, $W_+ \simeq T_+^{-1}$ and $W_- \simeq T_-^{-1}$ (probabilities for the right and for the left "jumps"). If $\langle \log(W_+) \rangle = \langle \log(W_-) \rangle$, the average displacement of the particle is zero, while the average square displacement behaves asymptotically as $\sim (\log t)^4$ [14,21]. This behavior is called as ultraslow (Sinai) diffusion, and may be regarded as a localization in the length scale of the ordinary diffusion [14]. If the external field is zero, $E = 0$, the ultraslow diffusion occurs in ratchets, which are symmetric in average.

To prove the existence of the ultraslow (Sinai) diffusion, we applied an approach commonly used to analyze this phenomenon [21]. Namely, we consider a motion of a particle in a segment of $N = 20$ ratchets with α distributed uniformly between 0.2 and 0.8, so that the average length of the left slope was equal to the average

length of the right slope. The reflecting left boundary and adsorbing right boundary were used. We calculated numerically the distribution $P(T_N)$ of times T_N , when the particle, initially located at the left boundary, reached the adsorbing boundary. For the ordinary 1d diffusion, the $T_N \gg 1$ asymptotics reads $P(T_N) \sim T_N^{-3/2}$, while for the Sinai diffusion $P(T_N) \sim T_N^{-1}$ [21]. The numerical results shown in Fig.3 illustrate that the ultraslow diffusion does take place in ratchets with the disorder.

The transition probability rates $W_{\pm}$, as well as the kinetic equation (6) refer to the time coarse-grained description and may be unambiguously applied only for $t \gg T_{\pm}$. To analyze the detailed balance or the thermodynamic flux $J(x) = \lim_{\Delta t \rightarrow 0} (N_+ - N_-)/\Delta t$, one needs more refined approach, since $\Delta t \ll T_{\pm}$. Here N_+ and N_- denotes correspondingly the number of particles that have crossed the plane located at x in positive and negative directions during the time interval Δt . To show that the asymmetry of the $T_{\pm}(x)$ does not imply the non-zero equilibrium flux, we write for the flux, e.g. at $x = L_+$ (see Fig.1): $J(x, t) = (1/\Delta t)[\int_{-L_-}^{L_+} P(x, t) \theta(\Delta t - T_+(x))dx - \int_{L_+}^{L_+ + L_0} P(x, t) \theta(\Delta t - T_-(x))dx]$, where the unit step function $\theta(t)$ guarantees that particles have crossed the plane within the Δt . Since $\Delta t \rightarrow 0$, we expand $T_{\pm}(x)$ [17] near $x = L_+$: $T_{\pm}(x) = |L_+ - x| e^{U_0/kT}/a_0 D$. As a result, a zero flux is obtained for the equilibrium probability density $P_{eq}(x)$.

In spite of a lack of the thermodynamic current at equilibrium, very unusual equilibrium fluctuations may be observed in circular ratchets. If particles may be labeled and if their paths may be traced, one observes that the average number of times that particle crosses some cross-section clockwise, $\langle S_+ \rangle$, is not equal at equilibrium to the average number of the counter-clockwise crossings, $\langle S_- \rangle$. Therefore, clockwise (or counter-clockwise) "drift of labels" takes place in equilibrium. Obviously, this is not a thermodynamical current, and no useful work may be extracted. We quantify the "drift of labels" by $S = (\langle S_+ \rangle / \langle S_- \rangle - 1)$. We also introduce the mean times of the clockwise turn over the ring, T_{C+} , and of the counter-clockwise turn, T_{C-} . If, again, $\exp(U_0/kT) \gg 1$ and $\tau_{\pm}/T_{\pm} \ll 1$, a straightforward solution of Eq.(3) gives for the ring of M ratchets:

$$T_{C\pm} = \frac{e^{U_0/kT}}{Da_0} \left(\frac{M(M-1)}{2a_0} + \frac{M}{a_{\pm}} \right) \quad (8)$$

and we estimate S as $(T_{C+}^{-1}/T_{C-}^{-1} - 1) = \frac{4\alpha-2}{(M+1)-2\alpha}$. The "flux of labels" disappears at large M , and thus it is an intrinsically microscopic phenomenon. Our numerical studies of the Brownian motion on the ring, of $M = 4$ ratchets with $\alpha = 0.8$ confirmed that the average current ceased to zero, while S relaxed to 0.29, which is in a good agreement with the theoretical estimate $S = 0.35$.

In conclusion, we studied the Brownian motion of particles in a ratchet potential under the internal white noise. We found that the inter-well mean-first passage

times have a right-left asymmetry for the equilibrium and near-equilibrium intrawell distributions of particles. We have shown analytically and numerically that this asymmetry gives rise to the following effects: (i) nonstationary transport; (ii) localization or ultraslow (Sinai) diffusion under disorder in the parameter of asymmetry of the ratchet potential; (iii) unusual equilibrium fluctuations in circular ratchets, which may be called as a "drift of labels".

We are indebted to M. Bier, R. Kapral, and A. Malevanets for valuable comments and discussions.

-
- [1] M.O.Magnasco, Phys.Rev.Lett. **71**, 1447 (1993).
 - [2] R.D.Astumian and M.Bier, Phys.Rev.Lett. **72**, 1766 (1994).
 - [3] C.R. Doering, W. Horsthemke and J. Riordan, Phys. Rev. Lett. **72**, 2984 (1994).
 - [4] M.O.Magnasco, Phys.Rev.Lett. **72**, 2656 (1994).
 - [5] M.M.Millonas, Phys.Rev.Lett. **74** 10, (1995).
 - [6] M.M.Millonas and D.R.Chialvo, Phys.Rev.Lett. **76**, 550 (1996).
 - [7] M.Bier, and R.D.Astumian, Phys.Rev.Lett. **76**, 4277 (1996); M.Bier, Phys. Lett. A **211**, 12 (1996); J.Rousselet, L.Salome, A.Ajdari, and J.Prost, Nature(London) **370**, 446 (1994); J.Prost, J.F.Chauwin, L.Peliti, and A.Ajdari, Phys.Rev.Lett. **72**, 2652 (1994)
 - [8] G.A.Cecchi and M.O.Magnasco, Phys.Rev.Lett. **76**, 1968 (1996).
 - [9] R.Bartussek, P.Reimann, and P.Hanggi, Phys.Rev.Lett. **76**, 1166 (1994)
 - [10] M.M.Millonas, and M.I.Dykman, Phys.Lett. **A185**, 65 (1994).
 - [11] J.Luczka, R.Bartussek, and P.Hanggi, Europhys. Lett. **31** 431 (1995); P.Jung, J.G.Kissner, and P.Hanggi, Phys.Rev.Lett. **76**, 3436 (1996).
 - [12] T.Hondo and Y.Sawada Phys.Rev.Lett. **75** 3269 (1995).
 - [13] Note, that the ensemble of the Brownian particles, being in equilibrium with the bath by their impulses, is not necessarily in a configurational equilibrium.
 - [14] J.P.Bouchaud and A.Georges, Phys.Reports, **195**, 127 (1990).
 - [15] M.Bier and R.D.Astumian, Phys.Rev.Lett. **71**, 1649 (1993); C.R.Doering and J.C.Gadoua, Phys.Rev.Lett. **69**, 2318 (1992).
 - [16] C.W.Gardiner, *Handbook of Stochastic Methods* (Springer, Berlin, 1985), 2nd ed.
 - [17] For $T_+(x)$ at $x > 0$ one has: $T_+(x > 0) = (e^{U_0/kT} - e^{a_+x/kT})/Da_+a_0 + (e^{-U_0/kT}e^{a_+x/kT} - 1)/Da_+a_-$, with similar expressions for $T_+(x < 0)$, and for $T_-(x)$.
 - [18] T.S.Elston, and C.R.Doering, J.Stat.Phys., **83**, 359 (1996).
 - [19] We found that the interwell transition rates were close to the reverse values of the MFPT for $U_0/kT \geq 0.1$.
 - [20] N.V.Brillantov, and A.I.Kviatkevitch, Sov. Phys. Solid State, v.31, 2060 (1989).

- [21] K.W.Kehr and K.P.N.Murthy, Phys.Rev. **A40**, 5728 (1990); K.P.N.Murthy, S.Rajasekar, and K.W.Kehr, J.Phys.A:Math.Gen. **27**, L107 (1994).

Fig. 1 Ratchet potential, $L_0 = L_+ + L_-$ (a); ratchet with a disorder of the asymmetry parameter (b).

Fig. 2 The average displacement versus time for the ensemble, initially localized at $x = 0$. Bottom insert: the

average velocity, V , versus the asymmetry parameter, α . Top insert: V versus the external force, E . The dots - simulations, the lines - theory; $kT = 0.085$, $\alpha = 0.8$, $\gamma = 1$, $L_0 = 1$, $U_0 = 1$.

Fig. 3 The distribution of escape times for the segment of 20 ratchets with reflecting left boundary. Bottom curve - the symmetric ratchet, $\alpha = 1/2$. Upper curve - the ratchet with a disorder; a flat distribution of α was used.





